

GET THIS ONE DONE.

If you're over 50, ask about PREVNAR 13[®] at your next visit to your doctor's office or pharmacy.

Learn more at AskAboutPREVNAR13.com

Please see Indications on page 1 and Important Safety Information on page 9 and full Prescribing Information for PREVNAR 13[®] inside pocket.



PREVNAR 13 is a registered trademark of Wyeth LLC.
Manufactured by Wyeth Pharmaceuticals Inc.



PP-PNA-USA-2357-01 ©2017 Pfizer Inc. All rights reserved.
Printed in the USA/April 2017

ADULTS 50+

When I protect myself,
I don't think this is scary.



Know what is scary
at my age?

**Pneumococcal
pneumonia.**

One dose of the PREVNAR 13[®] vaccine can help protect you.

You're over 50. Active. Young at heart. You take steps to protect your body when you're out doing things you love. And because your risk of getting pneumococcal pneumonia increases as you age, taking steps to help protect yourself from this potentially serious disease is just as important.



50 OR OLDER?


Ask about PREVNAR 13[®] at your next visit to your doctor's office or pharmacy.

INDICATIONS FOR PREVNAR 13[®]

- Prevnar 13[®] is a vaccine approved for adults 18 years of age and older for the prevention of pneumococcal pneumonia and invasive disease caused by 13 *Streptococcus pneumoniae* strains (1, 3, 4, 5, 6A, 6B, 7F, 9V, 14, 18C, 19A, 19F, and 23F)
- Prevnar 13[®] is not 100% effective and will only help protect against the 13 strains included in the vaccine

IMPORTANT SAFETY INFORMATION

- Prevnar 13[®] should not be given to anyone with a history of severe allergic reaction to any component of Prevnar 13[®] or any diphtheria toxoid-containing vaccine
- Adults with weakened immune systems (eg, HIV infection, leukemia) may have a reduced immune response




Help protect yourself from pneumococcal pneumonia.

GET THIS ONE DONE.

Pneumococcal pneumonia can seriously set you back.

Pneumococcal pneumonia is not a cold or the flu. It can be a serious lung infection that is caused by a common bacteria. It can be passed to others through a simple cough or a sneeze. It can even put you in the hospital and in severe cases, may lead to death.

But just one dose of the PREVNAR 13[®] vaccine can help prevent pneumococcal pneumonia. Get this one done. For you and those who depend on you.




If you're 50 or older,
your risk of **being hospitalized**
after getting pneumococcal pneumonia
is **8X higher than younger adults.**



50 OR OLDER?

Ask about PREVNAR 13[®]
at your next visit to your
doctor's office or pharmacy.

GET THIS ONE DONE.



Pneumococcal pneumonia can strike anywhere, anytime.

GET THIS ONE DONE.

Symptoms are distinct, can appear quickly, and may include:



Chest pain with difficulty breathing



A high fever, shaking chills



Excessive sweating



Fatigue



A cough with phlegm that persists or gets worse

Even if you're healthy, risk increases with age.

Chances are you go the extra mile to help prevent disease, like eating right and exercising. But as we get older, our immune system naturally weakens, making it harder for our bodies to fight off infections and disease. In fact, doctors say even healthy adults over 50 are at increased risk for pneumococcal pneumonia.

65 or older? The CDC recommends PREVNAR 13®.


The CDC recommends that adults 65 or older receive the PREVNAR 13® vaccine to help protect against pneumococcal disease. Immune response may be lower if PREVNAR 13® is given within one year after another pneumococcal vaccine.



Ask about PREVNAR 13® at your next visit to your doctor's office or pharmacy.

Information about PREVNAR 13® side effects.

The most common side effects reported in clinical trials were injection site reactions including pain, redness, and swelling at the injection site, and limitation of arm movement. Other commonly reported adverse reactions were fatigue, headache, muscle pain, joint pain, decreased appetite, vomiting, fever, chills, and rash. Adults with weakened immune systems (eg, HIV infection, leukemia) may have a reduced immune response. Please see additional Important Safety Information on page 9.



One dose of PREVNAR 13® can help protect you from pneumococcal pneumonia.

GET THIS ONE DONE.

IMPORTANT SAFETY INFORMATION

- Prevnar 13[®] should not be given to anyone with a history of severe allergic reaction to any component of Prevnar 13[®] or any diphtheria toxoid-containing vaccine
- Adults with weakened immune systems (eg, HIV infection, leukemia) may have a reduced immune response
- In adults, the most common side effects were pain, redness, and swelling at the injection site, limitation of arm movement, fatigue, headache, muscle pain, joint pain, decreased appetite, vomiting, fever, chills, and rash
- Ask your healthcare provider about the risks and benefits of Prevnar 13[®]. Only a healthcare provider can decide if Prevnar 13[®] is right for you

You are encouraged to report negative side effects of vaccines to the US Food and Drug Administration (FDA) and the Centers for Disease Control and Prevention (CDC). Visit www.vaers.hhs.gov or call 1-800-822-7967.

PREVNAR 13[®] may be available at no cost to you.

- If you're 65 or older, PREVNAR 13[®] may be covered by Medicare Part B with \$0 out-of-pocket cost.
- If you're between the ages of 50 and 64, check with your insurance provider.
- Need help paying for your Pfizer medicines? Pfizer Rx Pathways[™] is here to help. Learn more at PfizerRxPathways.com.



GET THIS ONE DONE.

Learn more at AskAboutPREVNAR13.com

Take one more step.

Help protect yourself from pneumococcal pneumonia.

GET THIS ONE DONE.

